AWARD NUMBER DAMD17-98-1-8108

TITLE: National Hospital Discharge Survey Data Analysis of Breast Cancer Between African American and White Women

PRINCIPAL INVESTIGATOR: KyungSook Kim, Ph.D.

CONTRACTING ORGANIZATION: Howard University

Washington, D.C. 20059

REPORT DATE: August 1999

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20001019 096

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

		<u> </u>		<u>-</u>
1. AGENCY USE ONLY (Leave blan	2. REPORT DATE August 1999	3. REPORT TYPE AND Annual Summary (2		
4. TITLE AND SUBTITLE National Hospital Discharge Sur American and White Women	vey Data Analysis of Breast Ca	ncer Between African		ng numbers 7-98-1-8108
6. AUTHOR(S)				
KyungSook Kim, Ph.	D.			
7. PERFORMING ORGANIZATION N Howard University Washington, D.C. 20059	AME(S) AND ADDRESS(ES)			RMING ORGANIZATION T NUMBER
9. SPONSORING / MONITORING ACU.S. Army Medical Research and Fort Detrick, Maryland 21702-5	nd Materiel Command	S)		SORING / MONITORING CY REPORT NUMBER
11. SUPPLEMENTARY NOTES	William Sales Westernach Browning			
12a. DISTRIBUTION / AVAILABILIT Approved for public release; dis			12b. DIST	TRIBUTION CODE
13. ABSTRACT (Maximum 200 wo	ords)		l	- Aller - Aller - Control
African American (AA) a 1987 data used for baseling between AA and white we women had proportionate rates for breast cancer base among AA than whites for higher among whites than combined, the mean (s.d.) stay ranged from 6.0 (10. (1.8) to 3.4 (2.1), and the the years 1988 and 1989.	ts to estimate age-specific hand white women through and the comparison. Breast cancer of the comparison of the various age of the various age of the various age of the various and the various and the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age. The various age of the various age of the various age. The various age of the various age of the various age. The various age of the various age of the various age. The various age of the various age of the various age. The various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the various age of the various age of the various age. The various age of the vari	nalyzing NHDS data cer admissions were e groups. For the 19 We analyzed age-spe OS data. In 1988, the ges 40-49. In 1989, ups below age 50. F 3) to 59.4 (14.8), the d.) number of diagn	from 19 not prop 987 data cific hos ose rates however for both e mean (oses ran	288 to 1994 with portionally equal young AA spital admission is were higher er, the rates were groups (s.d.) length of a ged from 3.0 to 1.6 (1.1) for
14. SUBJECT TERMS Breast Cancer, National Ho rate due to breast ca		, Hospital Admiss	sion	15. NUMBER OF PAGES 18 16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFI OF ABSTRACT Unclassified	CATION	20. LIMITATION OF ABSTRACT Unlimited

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.
Where copyrighted material is quoted, permission has been obtained to use such material.
Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.
Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.
In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).
For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.
In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.
In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.
In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

PI - Signature

8/16/99

Date

TABLE OF CONTENTS

		Page
Front Cover		1
Standard Form 29	98	2
Foreword		3
Table of Contents	s	4
Introducti	on	5
Body		5
Appendices		8
Figure 1.	Age-specific Hospital Admission rate of breast cancer by year	
	among White Women	9
Figure 2.	Age-specific Hospital Admission rate of breast cancer by year	
	among African American Women	10
Table 1.	Age-specific Hospital Admission rate of breast cancer by race:	
	Year 1987	11
Table 2.	Age-specific Hospital Admission rate of breast cancer by race:	
	Year 1988	12
Table 3.	Age-specific Hospital Admission rate of breast cancer by race:	
	Year 1989	13
Table 4.	Mean and standard deviation of age, length of stay, number of	
	diagnosis, and procedures by year and race	
	Discharge Status of breast cancer patients by year	
	Expected Source of payments of breast cancer patients by year	
	Marital Status of breast cancer patients by year	
Table 8.	Regional distribution of breast cancer patients by year	18

Introduction

The present study attempts to estimate age-specific hospital admission rates for breast cancer in African American (AA) and white women through analyzing NHDS data from 1988 to 1994 with 1987 data used for baseline comparison. Breast cancer admissions were not proportionally equal between AA and white women across the various age groups. For the 1987 data, young AA women had proportionately higher admission rates. It is necessary to see if this trend continued from 1988 to 1994, and if so, then specific guidelines may need to be set targeting this group of pre-menopausal AA women.

Body

We analyzed age-specific hospital admission rates for breast cancer based on 1988 and 1989 NHDS data. Tables 1-3 and Figures 1-2 show the distribution of female breast cancer cases and rates by age-group. In 1987, the relative risk of hospital admission due to breast cancer was higher among whites than AA for all age groups except at ages 40-49. In 1988, those rates were higher among AA than whites for all age groups except at ages 40-49. In 1989, however, the rates were higher among whites than those of AA in all age groups below age 50. In 1988 and 1989, there were 240,503 cases and 225,820 cases estimated in white women and 41,445 cases and 31,130 cases estimated in AA women, respectively. For both groups combined, the mean (s.d.) age ranged from 59.1 (14.3) to 59.4 (14.8), the mean (s.d.) length of a stay ranged from 6.0 (10.2) to 6.7 (8.8), the mean (s.d.) number of diagnoses ranged from 3.0 (1.8) to 3.4 (2.1), and the mean (s.d.) number of procedure ranged from 1.3 (1.0) to 1.6 (1.1) for the years 1988 and 1989 (Table 4).

Table 5 shows the discharge status of breast cancer patients from 1987 to 1989 for whites

and AA. The average percentage of routine discharge was higher among whites than AA. However, the average percentage of death was lower among whites than AA. Table 6 shows the percentages of expected sources of payments by year and race. While the proportion of Medicare patients constituted about 37% among whites, it was about 28% among AA for the year 1987 and 1988 but sharply increased to 42% at 1989. AA women were four times more likely to be paid by Medicaid than whites. The married category had the highest proportion in 1987 while the unknown category had the highest proportion in 1989 among whites. For AA women, single/divorce/widow/separate category was the highest for the years 1987 to 1989 (Table 7). The Southern regions had the highest rates, while the West had the lowest rates of hospital admission due to breast cancer. These rates were the same for both whites and AA for the year 1987 to 1989 (Table 8).

From RIDIT analysis, we found that mean ridit was 0.495 in 1988 and 0.504 in 1989 among whites with year 1987 as a reference group. That means, the subjects in 1988 were slightly younger than the subjects in 1987. Among whites, the subjects age distribution was slightly older in 1989 than subjects in 1987. However, among AA, mean ridit was 0.607 in 1988 and 0.609 in 1989 with year 1987 as a reference group. That is, the subjects in 1988 and in 1989 were significantly older than the subjects in 1987.

Figure 1 revealed that in 1987 the hospital admission rate due to breast cancer among whites increased linearly until age group 60-69 where after it began to plateau. Comparing age-specific rates of 1988 and 1989 with those of 1987, all the rates were lower than those of 1987 and the gap was greatest among the age group 60-69 for both years. The rate in the age group 70-84 was decreased in 1989 when compared to that of 1988. Figure 2 revealed that among AA, there was a notably low hospital admission rate among the 50-59 and 70-84 age groups in 1987.


The rates increased sharply in these age groups in 1988. In 1989, the rates demonstrated a more gradual increase until the age group 60-69 after which the rates increased sharply. With additional analysis of the 1990 through 1994 NHDS data, long term trends may be identified.

Difficulties:

Unfortunately we have not been able to identify and hire a statistician/research associate as of yet.

If we cannot hire a person soon, we may need to seek an extension of the study. Due to the vacancy of the position, analysis of the comorbid diagnoses and procedures performed between these two groups for all the study periods has been delayed.

APPENDICES



rate: per 100,000

Figure 1. Age-specific Hospital Admission rate of breast cancer by year among White Women




Figure 2. Age-specific Hospital Admission rate of breast cancer by year among African-American Women

Table 1. Age-specific Hospital Admission rate of Breast cancer by race: year 1987

	White				<u>AA</u>		
Age	Rate *	n	%	Rate *	n	%	RR**
20-39	76.89	26,233	9.9	74.63	4,025	15.9	0.97
40-49	351.61	42,753	16.1	433.67	6,813	26.9	1.23
50-59	566.1	55,688	20.9	257.6	3,220	12.7	0.46
60-69	740.5	72,478	27.2	712.4	7,295	28.8	0.96
70-84	774.5	69,037	25.9	509.4	3,963	15.7	0.66
		266,189			25,316		

^{*} rate per 100,000

** Relative Risk: AA rate/ White rate

Table 2. Age-specific Hospital Admission rate of Breast cancer by race: year 1988

	<u>White</u>				<u>AA</u>		
Age	Rate *	n	%	Rate *	n	%	RR**
20-39	69.3	23,562	9.8	79.5	4,333	10.5	1.15
40-49	340.3	43,270	18.0	252.3	4,135	10.0	0.74
50-59	538.1	52,923	22.0	815.6	10,350	25.0	1.52
60-69	540.1	53,796	22.4	992.4	10,569	25.5	1.84
70-84	710.2	66,952	27.7	1440.6	12,058	29.1	2.03
 		240,503			41,445		

^{*} rate per 100,000

** Relative Risk: AA rate/ White rate

Table 3. Age-specific Hospital Admission rate of Breast cancer by race: year 1989

	White				<u>AA</u>		
Age	Rate *	n	%	Rate *	n	%	RR**
20-39	49.9	16,927	7.5	46.3	2,541	8.2	0.93
40-49	317.0	41,942	18.6	312.1	5,347	17.2	0.99
50-59	500.7	49,207	21.8	502.9	6,472	20.8	1.00
60-69	555.5	55,382	24.5	599.0	6,481	20.8	1.08
70-84	651.2	62,362	27.6	1210.5	10,289	33.0	1.86
	- ***	225,820			31,130		

^{*} rate per 100,000

** Relative Risk: AA rate/ White rate

Table 4.Mean and standard deviation of age, length of stay, number of diagnosis, and procedures by year and race

	White					
	<u> 1987</u>		<u> 1988</u>		<u> 1989</u>	
	⋝	s.d.	⋝	s.d.	⋝	s.d.
	50.00	10.07	50.06	140	50.44	12.0
Age	58.92	13.87	59.06	14.3	59.44	13.8
Length of stay	6.07	7.26	6.13	7.6	5.97	10.2
Number of diagnosis	3.01	1.92	3.04	1.8	3.20	1.87
Number of procedures	1.57	1.09	1.58	1.1	1.61	1.12

	<u>1987</u> ⊼	s.d.	<u>1988</u> ⊼	s.d.	<u>1989</u> ⊼	s.d.
Age	53.55	14.69	59.44	12.8	59.44	14.8
Length of stay	7.59	10.3	6.27	8.0	6.72	8.8
Number of diagnosis	3.09	1.95	3.01	1.77	3.38	2.06
Number of procedures	1.71	1.01	1.49	1.0	1.27	0.97

Table 5. Discharge Status of breast cancer patients by year

	<u>1987</u> n	%	White 1988 n	%	<u>1989</u> n	%
Routine discharge	234,777	88.1	212,709	88.4	199,064	88.2
Transfer to short- term hospital	1,519	0.6	1,198	0.5	1,979	0.9
Transfer to long- term care	9,244	3.5	7,293	3.0	8,520	3.8
Dead	10,520	4.0	11,607	4.8	7,395	3.3
Not stated	10,110	3.8	7,696	3.2	7,699	3.4
		Africa	an American			
	<u>1987</u> n	%	<u>1988</u> n	%	<u>1989</u> n	%
Routine discharge	21,147	83.5	37,014	89.3	26,949	86.6
Transfer to short- term hospital	205	0.8	0	0.0	45	0.1
Transfer to long- term care	609	2.4	453	1.1	1,447	4.7
Dead	2,656	10.5	1,781	4.3	1,926	6.2
Not stated	575	2.3	1,875	4.5	763	2.5

Table 6. Expected Source of payments of breast cancer patients by year

	<u>1987</u>	%	White 1988 n	%	<u>1989</u>	%
Medicare	103,099	38.7	88,836	36.9	85,324	37.8
Medicaid	8,104	3.0	6,038	2.5	7,519	3.3
Blue Cross/ other private ins.	140,364	52.7	129,869	54.0	108,148	47.9

	<u>1987</u> n	%	1988 n	%	1989 n	%
Medicare	7,075	28.0	11,769	28.4	13,157	42.3
Medicaid	2,971	11.7	6,588	15.9	3,851	12.4
Blue Cross/ other private ins.	12,339	48.7	18,526	44.7	11,645	37.4

Table 7. Marital Status of breast cancer patients by year

·	<u>1987</u>	%	White 1988 n	%	<u>1989</u>	%
Married	125,137	47	87,269	36.3	67,639	30.0
Single/Widow/ Divorce/Separate	77,845	29.2	45,736	19.0	51,535	22.8
Unknown	63,207	23.7	107,498	44.7	106,646	47.2

	<u>1987</u>	%	<u>198</u>	<u>8</u> %	<u>1989</u>	%
Married	6,658	26.3	13,915	33.6	7,503	24.1
Single/Widow/ Divorce/Separate	11,972	47.3	15,293	36.9	15,333	49.3
Unknown	6,686	26.4	12,237	29.5	8,294	26.6

Table 8 . Regional distribution of breast cancer patients by year

	1987		White 1988		1989	
	n	%	n	%	n	%
North East	58,315	21.9	72,306	30.1	64,731	28.7
Mid West	76,671	28.8	51,085	21.2	64,295	28.5
South	82,310	30.9	83,941	34.9	76,604	33.9
West	48,893	18.4	33,171	13.8	20,190	8.9

	<u>1987</u>	%	<u>1988</u>	%	<u>1989</u>	%
North East	6,429	25.4	13,912	33.6	10,064	32.3
Mid West	4,737	18.7	7,370	17.8	5,658	18.2
South	12,912	51.0	18,858	45.5	13,910	44.7
West	1,238	4.9	1,305	3.1	1,498	4.8